



ARRAY

3X

## WHEAT/GLUTEN PROTEOME REACTIVITY & AUTOIMMUNITY™



2 mL Serum

Antigens Tested (IgG and IgA tested separately for each antigen)

- Wheat
- Wheat Germ Agglutinin
- Non-Gluten Proteins-A\*
- Non-Gluten Proteins-B\*
- Gliadin Toxic Peptides\*
- Native + Deamidated Alpha-Gliadin-33-mer
- Alpha-Gliadin-17-mer
- Gamma-Gliadin-15-mer
- Omega-Gliadin-17-mer
- Glutenin-21-mer
- Gluteomorphin+Prodynorphin
- Gliadin-Transglutaminase Complex
- Microbial Transglutaminase\*
- Transglutaminase-2
- Transglutaminase-3
- Transglutaminase-6

\* Newly added antigens

### CLINICAL USE:

- Identify possible Celiac disease, non-celiac gluten sensitivity, dermatitis herpetiformis, gluten ataxia or other wheat/gluten-related disorder.
- Assess autoimmune reactivity associated with wheat proteins and peptides.

### RECOMMENDED FOR PATIENTS WHO:

- Have non-responsive GI symptoms.
- Present multiple-symptom complaints (including Chronic Fatigue Syndrome and Fibromyalgia).
- Suffer from depression or neuro-autoimmunity.

 **CYREX**<sup>®</sup>  
Laboratories  
[www.JoinCyrex.com](http://www.JoinCyrex.com)

# WHEAT PROTEINS • PEPTIDES

